

NONDIFFERENTIABLE GENERALIZED MINIMAX FRACTIONAL PROGRAMMING UNDER (Φ, ρ) -INVEXITY

B.B. UPADHYAY

*Department of Mathematics, Indian Institute of Technology Patna,
Patna-801103, India
bhooshan@iitp.ac.in*

T. ANTCZAK

*Faculty of Mathematics and Computer Science, University of Lodz', Banacha 22,
90-238 Lodz', Poland
antczak@math.uni.lodz.pl*

S.K. MISHRA

*Department of Mathematics, Institute of Science, Banaras Hindu University,
Varanasi-221005, India
bhu.skmishra@gmail.com*

K. SHUKLA

*Department of Mathematics, GLA University, Mathura, Mathura-281406, India
bhu.kalpanabhu@gmail.com*

Received: September 2020 / Accepted: June 2021

Abstract: In this paper, a class of nonconvex nondifferentiable generalized minimax fractional programming problems is considered. Sufficient optimality conditions for the considered nondifferentiable generalized minimax fractional programming problem are established under the concept of (Φ, ρ) -invexity. Further, two types of dual models are formulated and various duality theorems relating to the primal minimax fractional programming problem and dual problems are established. The results established in the paper generalize and extend several known results in the literature to a wider class of nondifferentiable minimax fractional programming problems. To the best of our knowledge, these results have not been established till now.

Keywords: Nondifferentiable Minimax Fractional Problem, Duality, (Φ, ρ) -invexity,

Sufficient Conditions.

MSC: 90C26, 90C32, 49K35.

1. INTRODUCTION

Minimax programming has been an interesting field of active research for a long time. These problems are of pivotal importance in many areas of modern research such as economics, engineering design, portfolio selection, game theory, rational Chebyshev approximations and financial planning, see [7, 8, 43] and the references cited therein. Necessary optimality conditions for finite-dimensional constrained minimax problems in terms of Lagrange multipliers have been originally investigated by Bram [11] and Danskin [15]. Schmitendorf [42] has established the necessary and sufficient optimality conditions for the following minimax programming problem:

$$(P) \quad \min \sup_{y \in Y} f(x, y)$$

subject to $h(x) \leq 0$,

where $f(., .) : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ and $h(.) : \mathbb{R}^n \rightarrow \mathbb{R}^p$ are differentiable convex functions and Y is a compact subset of \mathbb{R}^m . Bector and Bhatia [9] and Weir [48] relaxed the convexity assumption in proving the sufficient optimality conditions for problem (P) and formulated several dual models. They proved such results under pseudoconvexity and quasiconvexity assumptions imposed on the functions constituting problem (P) and its duals. Bector et al. [10] derived duality results for minimax programming problems involving V -invex functions.

The concept of invex functions introduced by Hanson [24] and named by Craven [14] is a significant generalization of convex functions. The theory of mathematical programming has grown remarkably, when further extensions of invexity have been introduced to establish the optimality conditions and duality results. Preda [41] introduced the concept of (F, ρ) -convexity as an extension of F -convexity defined by Hanson and Mond [25], whereas the concept of ρ -convexity was introduced by Vial [46]. Jeyakumar [28] generalized the Vial [46] notion of ρ -convexity to ρ -invexity. In a recent work, Yuan et al. [50] introduced a unified formulation of generalized convexity, called (C, α, ρ, d) -convexity. The (C, α, ρ, d) -convexity extends the (F, α, ρ, d) -convexity introduced by Liang et al. [31] by relaxing the sublinearity of the scale function to convexity. Caristi et al. [12] introduced the notion of (Φ, ρ) -invexity for differentiable scalar optimization problems, which generalizes invexity as well as (F, ρ) -convexity. Recently, Antczak and Stasiak [6] generalized the concept of (Φ, ρ) -invexity to a nondifferentiable case and they introduced the definition of a locally Lipschitz (Φ, ρ) -invex function. Very recently, nonsmooth semi-infinite minimax programming problems with locally Lipschitz (Φ, ρ) -invex and generalized locally Lipschitz (Φ, ρ) -invex functions have been studied by Liu et al. [34] and Upadhyay and Mishra [45].

Many authors investigated the optimality conditions and duality results for minimax fractional programming problems using generalized convexity assumptions, see for example [1, 3, 4, 5, 13, 16, 17, 18, 19, 20, 21, 32, 33, 34, 38, 39, 44, 45] and the references cited therein. Lai et al. [30] established the necessary and sufficient optimality conditions and Lai and Lee [29] obtained duality results for a class of nondifferentiable minimax programming problems with generalized convex functions. Several authors have developed interesting results in nondifferentiable minimax fractional programming problems; see for example [26, 27, 33, 36, 37, 40] and the references therein. Recently, the concept of nondifferentiable (Φ, ρ) -invexity was used by Antczak [2] in proving the sufficient optimality conditions and various duality results for a class of nondifferentiable minimax programming problems.

Motivated by the works of Caristi et al. [12], Lai and Lee [29], Lai et al. [30], Upadhyay and Mishra [45] and Yuan et al. [50], we consider a class of nonconvex nondifferentiable generalized minimax fractional programming problems with (Φ, ρ) -invex functions. We establish the sufficient optimality conditions for such a nondifferentiable optimization problem. The necessary optimality criteria are then used to formulate two types of dual models for the considered nondifferentiable generalized minimax programming problem. Further, under (Φ, ρ) -invexity hypotheses, weak, strong and strict converse duality results are established between the considered nondifferentiable generalized minimax programming problem and its duals formulated in the paper.

2. DEFINITIONS and PRELIMINARES

Let \mathbb{R}^n be the n -dimensional Euclidean space and \mathbb{R}_+^n be the non-negative orthant of \mathbb{R}^n . Let $\emptyset \neq X_0 \subseteq \mathbb{R}^n$ and $\langle \cdot, \cdot \rangle$ denotes the Euclidean inner product.

We consider the following nondifferentiable generalized minimax fractional programming problem:

$$(P) \quad \inf_{x \in \mathbb{R}^n} \sup_{y \in Y} \left\{ \phi(x, y) := \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} \right\}$$

$$\text{subject to } h_j(x) \leq 0, j \in J := \{1, \dots, p\},$$

where Y is a compact subset of \mathbb{R}^m , $f(\cdot, \cdot), g(\cdot, \cdot) : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ and $h : \mathbb{R}^n \rightarrow \mathbb{R}^p$ are C^1 -mappings. Let A and B be $n \times n$ positive semidefinite matrices. The problem (P) is a nondifferentiable optimization problem if either A or B is nonzero. If A and B are null matrices, problem (P) is a differentiable generalized minimax fractional programming problem.

Let $X := \{x \in \mathbb{R}^n \mid h_j(x) \leq 0, j \in J\}$ be the set of all feasible solutions for (P). Assume that for each $(x, y) \in X \times Y$, $f(x, y) + \langle x, Ax \rangle^{1/2} \geq 0$ and $g(x, y) - \langle x, Bx \rangle^{1/2} > 0$. Let us define the following sets for every $x \in X$:

$$J(x) := \{j \in J \mid h_j(x) = 0\},$$

$$Y(x) := \left\{ y \in Y \mid \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} = \sup_{z \in Y} \frac{f(x, z) + \langle x, Ax \rangle^{1/2}}{g(x, z) - \langle x, Bx \rangle^{1/2}} \right\},$$

$$K(x) := \left\{ (s, t, \bar{y}) \in N \times \mathbb{R}_+^s \times \mathbb{R}^{ms} : 1 \leq s \leq n+1, t = (t_1, \dots, t_s) \in \mathbb{R}_+^s \right.$$

$$\left. \text{with } \sum_{i=1}^s t_i = 1 \text{ and } \bar{y} = (\bar{y}_1, \dots, \bar{y}_m) \text{ with } \bar{y}_i \in Y(x), i = 1, \dots, s \right\}.$$

Since f and g are continuously differentiable and Y is a compact subset of \mathbb{R}^m , it follows that, for each $x^* \in X$, $Y(x^*) \neq \emptyset$ and, for any $\bar{y}_i \in Y(x^*)$, we can find a positive constant k_0 such that $k_0 = \phi(x^*, \bar{y}_i)$. The following generalized Schwartz inequality will be needed in our considerations:

$$\langle x, A\nu \rangle \leq \langle x, Ax \rangle^{1/2} \langle \nu, A\nu \rangle^{1/2}, \forall x, \nu \in \mathbb{R}^n. \quad (1)$$

The equality holds if $Ax = \lambda A\nu$, for some $\lambda \geq 0$.

Hence, if $\langle \nu, A\nu \rangle^{1/2} \leq 1$, we have

$$\langle x, A\nu \rangle \leq \langle x, Ax \rangle^{1/2}. \quad (2)$$

Now on, we assume that an element of $(n+1)$ -dimensional Euclidean space \mathbb{R}^{n+1} is represented as the ordered pair (y, r) with $y \in \mathbb{R}^n$ and $r \in \mathbb{R}$. Let $\Phi : X_0 \times X_0 \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ and ρ be a real number such that $\Phi(x, u, \cdot)$ is convex on \mathbb{R}^{n+1} and $\Phi(x, u, (0, r)) \geq 0$, for all $(x, u) \in X_0 \times X_0$ and $r \geq 0$.

In order to relax the convexity assumptions, we use the following definitions from Ferrara and Stefanescu [24].

Definition 1. A differentiable function $f : X_0 \rightarrow \mathbb{R}$ is said to be (Φ, ρ) -invex at $u \in X_0$ on X , if for all $x \in X$, we have

$$f(x) - f(u) \geq \Phi(x, u, (\nabla f(u), \rho)).$$

The following example is taken from Upadhyay and Mishra [45]

Example 2. Consider the function $f : D \rightarrow \mathbb{R}$, given by

$$f(x) = (x_1 + 1)(x_2 + 2),$$

where $D =]-1, 1[\times]-1, 1[$. Define

$$\Phi(x, y, (z, \rho)) = 2(2^\rho - 1) |(x_1 - y_1)(x_2 - y_2)| + \langle z, x - y \rangle.$$

The function f is (Φ, ρ) -invex at $y = 0$ for $\rho = \frac{1}{2}$ on X , where $X = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 x_2 \geq 0\}$.

Definition 3. A differentiable function $f : X_0 \rightarrow \mathbb{R}$ is said to be strictly (Φ, ρ) -invex at $u \in X_0$ on X , if for all $x \in X$, we have

$$f(x) - f(u) > \Phi(x, u, (\nabla f(u), \rho)).$$

Remark 4. [12] If f_1 is (Φ, ρ_1) -invex and f_2 is (Φ, ρ_2) -invex, then $\lambda f_1 + (1 - \lambda)f_2$ is $(\Phi, \lambda\rho_1 + (1 - \lambda)\rho_2)$ -invex, whenever $\lambda \in [0, 1]$.

Remark 5. The notion of (Φ, ρ) -invexity generalizes and extends a number of generalized convexity notions. Indeed, from Definition 1, there are the following special cases:

(1) If $\Phi(x, u, (\nabla f(u), \rho)) = \langle \nabla f(u), \eta(x, u) \rangle$, where $\eta : X_0 \times X_0 \rightarrow \mathbb{R}^n$, then (Φ, ρ) -invexity reduces to the definition of invex function (with respect to η) introduced by Hanson [24].

(2) If $\Phi(x, u, (\nabla f(u), \rho)) = \frac{1}{b(x, u)} \langle \nabla f(u), \eta(x, u) \rangle$, where $b : X_0 \times X_0 \rightarrow \mathbb{R}_+ / 0$, and $\eta : X_0 \times X_0 \rightarrow \mathbb{R}^n$, then (Φ, ρ) -invexity reduces to the definition of b -invex function (with respect to η) (see, Mishra et al. [40]).

(3) If $\Phi(x, u, (\nabla f(u), \rho)) = F(x, u, \nabla f(u))$, where $F(x, u, \cdot)$ is a sublinear function in the third argument, then the (Φ, ρ) -invexity reduces to the definition of F -convexity introduced by Hanson and Mond [25].

(4) If $\Phi(x, u, (\nabla f(u), \rho)) = \langle \nabla f(u), (x - u) \rangle + \rho \|x - u\|^2$, then (Φ, ρ) -invexity reduces to the definition of ρ -convex function introduced by Vial [46].

(5) If $\Phi(x, u, (\nabla f(u), \rho)) = \langle \nabla f(u), \eta(x, u) \rangle + \rho \|d(x, u)\|^2$, where $\eta, d : X_0 \times X_0 \rightarrow \mathbb{R}^n$ and $d(x, y) \neq 0$, whenever $x \neq y$, then (Φ, ρ) -invexity reduces to the definition of ρ -invex function (with respect to η and d) introduced by Jeyakumar [28].

(6) If $\Phi(x, u, (\nabla f(u), \rho)) = F(x, u, \nabla f(u)) + \rho \|d(x, u)\|^2$, where $F(x, u, \cdot)$ is a sublinear functional in the third argument, then the (Φ, ρ) -invexity reduces to the definition of (Φ, ρ) -convexity introduced by Preda [41].

(7) Let $C : X_0 \times X_0 \times \mathbb{R}^n \rightarrow \mathbb{R}$, $\alpha : X_0 \times X_0 \rightarrow \mathbb{R}_+ \setminus \{0\}$ and $d : X_0 \times X_0 \rightarrow \mathbb{R}_+$ be the functions such that C is convex in the third argument satisfying $C(x, a, 0) = 0$, for all $(x, a) \in X_0 \times X_0$ and $d(x, a) = 0$ if and only if $x = a$. If $\Phi(x, a, (y, \rho)) = C(x, a, y)\alpha(x, a) + \rho d(x, a)$ and ρ is a constant, then the definition of a (Φ, ρ) -invex function reduces to the definition of (C, α, ρ, d) -convexity introduced by Yuan et al. [50].

3. OPTIMALITY CONDITIONS

In this section, we establish the sufficient optimality conditions for the problem (P) under (Φ, ρ) -invexity assumptions.

Before establishing the sufficient optimality conditions, we state the following necessary optimality conditions for the problem (P) established in Lai et al. [30]

Theorem 6. (Necessary optimality conditions) Let x^* be an optimal solution for (P) satisfying $\langle x^*, Ax^* \rangle > 0$, $\langle x^*, Bx^* \rangle > 0$ and $\nabla h_j(x^*), j \in J(x^*)$ are linearly

independent. Then, there exist $(s, t^*, \bar{y}) \in K(x^*)$, $k_0 \in \mathbb{R}_+$, $u, \nu \in \mathbb{R}^n$ and $\mu^* \in \mathbb{R}_+^p$, such that

$$\sum_{i=1}^s t_i^* \{ \nabla f(x^*, \bar{y}_i) + Au - k_0 (\nabla g(x^*, \bar{y}_i) - B\nu) \} + \nabla \langle \mu^*, h(x^*) \rangle = 0, \quad (3)$$

$$f(x^*, \bar{y}_i) + \langle x^*, Ax^* \rangle^{1/2} - k_0 \left(g(x^*, \bar{y}_i) - \langle x^*, Bx^* \rangle^{1/2} \right) = 0, i = 1, \dots, s, \quad (4)$$

$$\langle \mu^*, h(x^*) \rangle = 0, \quad (5)$$

$$t_i^* \geq 0, i = 1, \dots, s, \text{ with } \sum_{i=1}^s t_i^* = 1, \quad (6)$$

$$\langle u, Au \rangle \leq 1, \langle \nu, B\nu \rangle \leq 1, \quad (7)$$

$$\langle x^*, Au \rangle = \langle x^*, Ax^* \rangle^{1/2}, \langle x^*, B\nu \rangle = \langle x^*, Bx^* \rangle^{1/2}. \quad (8)$$

In the above theorem, we observe that both the matrices A and B are positive definite at the solution x^* . If one of $\langle x^*, Ax^* \rangle$ and $\langle x^*, Bx^* \rangle$ is zero or both A and B are singular at x^* , then, for $s, t^*, \bar{y} \in K(x^*)$, we can take a set $Z_{\bar{y}}(x^*)$ as defined in Lai et al. [30] by

$$Z_{\bar{y}}(x^*) := \{z \in \mathbb{R}^n \mid \langle \nabla h_j(x^*), z \rangle \leq 0, j \in J(x^*)\}$$

satisfying any one of the following three assumptions:

$$(i) \langle x^*, Ax^* \rangle > 0, \langle x^*, Bx^* \rangle = 0$$

$$\Rightarrow \left\langle \sum_{i=1}^s t_i^* \nabla f(x^*, \bar{y}_i) + \frac{Ax^*}{\langle Ax^*, x^* \rangle^{1/2}} - k_0 \nabla g(x^*, \bar{y}_i), z \right\rangle + \langle (k_0^2 B)z, z \rangle^{1/2} < 0,$$

$$(ii) \langle x^*, Ax^* \rangle = 0, \langle x^*, Bx^* \rangle > 0$$

$$\Rightarrow \left\langle \sum_{i=1}^s t_i^* \left(\nabla f(x^*, \bar{y}_i) - k_0 \left(\nabla g(x^*, \bar{y}_i) - \frac{Bx^*}{\langle Bx^*, x^* \rangle^{1/2}} \right) \right), z \right\rangle + \langle Bz, z \rangle^{1/2} < 0,$$

$$(iii) \langle x^*, Ax^* \rangle = 0, \langle x^*, Bx^* \rangle = 0$$

$$\Rightarrow \left\langle \sum_{i=1}^s t_i^* \nabla f(x^*, \bar{y}_i) - k_0 \nabla g(x^*, \bar{y}_i), z \right\rangle + \langle (k_0 B)z, z \rangle^{1/2} + \langle Bz, z \rangle^{1/2} < 0.$$

If we assume that $Z_{\bar{y}}(x^*) = \emptyset$ in Theorem 6, then the result of Theorem 6 still holds.

In the next theorem, we denote

$$\phi_0(\cdot) := \sum_{i=1}^s t_i^* \{ (f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle) - k_0 (g(\cdot, \bar{y}_i) - \langle \cdot, B\nu \rangle) \}.$$

Theorem 7. (Sufficient optimality conditions) Let $x^* \in X$ be a feasible solution for (P) and there exist $k_0 \in \mathbb{R}_+$, $(s, t^*, \bar{y}) \in K(x^*)$, $u, \nu \in \mathbb{R}^n$ and $\mu^* \in \mathbb{R}_+^p$ satisfying (3)-(8). Further, assume that $f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle$ and $-g(\cdot, \bar{y}_i) + \langle \cdot, B\nu \rangle$, $i = 1, \dots, s$, are (Φ, ρ_i) -invex and $(\Phi, \bar{\rho}_i)$ -invex at x^* on X , respectively, and $h_j(\cdot)$, $j = 1, \dots, p$, is (Φ, ρ_j^*) -invex at x^* on X such that, the inequality

$$\sum_{i=1}^s t_i^* (\rho_i + k_0 \bar{\rho}_i) + \sum_{j=1}^p \mu_j^* \rho_j^* \geq 0 \quad (9)$$

holds. Then x^* is an optimal solution for (P).

Proof. We proceed by contradiction. Suppose that x^* is not an optimal solution for (P). Then, there exists $\tilde{x} \in X$, such that

$$\sup_{y \in Y} \frac{f(\tilde{x}, y) + \langle \tilde{x}, A\tilde{x} \rangle^{1/2}}{g(\tilde{x}, y) - \langle \tilde{x}, B\tilde{x} \rangle^{1/2}} < \sup_{y \in Y} \frac{f(x^*, y) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, y) - \langle x^*, Bx^* \rangle^{1/2}}.$$

We observe that

$$\sup_{y \in Y} \frac{f(x^*, y) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, y) - \langle x^*, Bx^* \rangle^{1/2}} = \frac{f(x^*, \bar{y}_i) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, \bar{y}_i) - \langle x^*, Bx^* \rangle^{1/2}} = k_0,$$

for $\bar{y}_i \in Y(x^*)$, $i = 1, \dots, s$ and

$$\frac{f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, A\tilde{x} \rangle^{1/2}}{g(\tilde{x}, \bar{y}_i) - \langle \tilde{x}, B\tilde{x} \rangle^{1/2}} \leq \sup_{y \in Y} \frac{f(\tilde{x}, y) + \langle \tilde{x}, A\tilde{x} \rangle^{1/2}}{g(\tilde{x}, y) - \langle \tilde{x}, B\tilde{x} \rangle^{1/2}}.$$

Thus, we have

$$\frac{f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, A\tilde{x} \rangle^{1/2}}{g(\tilde{x}, \bar{y}_i) - \langle \tilde{x}, B\tilde{x} \rangle^{1/2}} < k_0, i = 1, \dots, s.$$

The above inequalities yield

$$f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, A\tilde{x} \rangle^{1/2} - k_0 (g(\tilde{x}, \bar{y}_i) - \langle \tilde{x}, B\tilde{x} \rangle^{1/2}) < 0, i = 1, \dots, s. \quad (10)$$

From (1), (4), (6), (7) – (8) and (10), we get

$$\begin{aligned} \phi_0(\tilde{x}) &= \sum_{i=1}^s t_i^* \left\{ \left(f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, Au \rangle^{1/2} \right) - k_0 \left(g(\tilde{x}, \bar{y}_i) - \langle \tilde{x}, B\nu \rangle^{1/2} \right) \right\} \\ &\leq \sum_{i=1}^s t_i^* \left\{ \left(f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, A\tilde{x} \rangle^{1/2} \right) - k_0 \left(g(\tilde{x}, \bar{y}_i) - \langle \tilde{x}, B\tilde{x} \rangle^{1/2} \right) \right\} \\ &< 0 = \sum_{i=1}^s t_i^* \left\{ \left(f(x^*, \bar{y}_i) + \langle x^*, Ax^* \rangle^{1/2} \right) - k_0 \left(g(x^*, \bar{y}_i) - \langle x^*, Bx^* \rangle^{1/2} \right) \right\} \\ &= \sum_{i=1}^s t_i^* \left\{ \left(f(x^*, \bar{y}_i) + \langle x^*, Au \rangle \right) - k_0 \left(g(x^*, \bar{y}_i) - \langle x^*, B\nu \rangle \right) \right\} = \phi_0(x^*). \end{aligned}$$

Thus,

$$\begin{aligned} & \sum_{i=1}^s t_i^* \{ (f(x, \bar{y}_i) + \langle x, Au \rangle) - (f(x^*, \bar{y}_i) + \langle x^*, Au \rangle) \} \\ & + \sum_{i=1}^s t_i^* k_0 \{ - (g(x, \bar{y}_i) - \langle x, Bv \rangle) + (g(x^*, \bar{y}_i) - \langle x^*, Bv \rangle) \} < 0. \end{aligned}$$

By (5) and $x \in X$, it follows that

$$\begin{aligned} & \sum_{i=1}^s t_i^* \{ (f(x, \bar{y}_i) + \langle x, Au \rangle) - (f(x^*, \bar{y}_i) + \langle x^*, Au \rangle) \} + \sum_{i=1}^s t_i^* k_0 \{ - (g(x, \bar{y}_i) - \\ & \langle x, Bv \rangle) + (g(x^*, \bar{y}_i) - \langle x^*, Bv \rangle) \} + \sum_{j=1}^p \mu_j^* (h_j(x) - h_j(x^*)) < 0. \quad (11) \end{aligned}$$

By the (Φ, ρ_i) -invexity of $(f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle)$ and $(\Phi, \bar{\rho}_i)$ -invexity of $(g(\cdot, \bar{y}_i) - \langle \cdot, Bv \rangle)$ at x^* , for $i = 1, \dots, s$, and for all $\tilde{x} \in X$, it follows that

$$\begin{aligned} & f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, Au \rangle - f(x^*, \bar{y}_i) - \langle x^*, Au \rangle \geq \\ & \Phi(\tilde{x}, x^*, (\nabla f(x^*, \bar{y}_i) + Au, \rho_i)), i = 1, \dots, s \end{aligned} \quad (12)$$

and

$$\begin{aligned} & -g(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, Bv \rangle + g(x^*, \bar{y}_i) - \langle x^*, Bv \rangle \geq \\ & \Phi(\tilde{x}, x^*, (\nabla f(x^*, \bar{y}_i) + Bv, \bar{\rho}_i)), i = 1, \dots, s. \end{aligned} \quad (13)$$

The (Φ, ρ_j^*) -invexity of $h_j(\cdot)$ at x^* implies that

$$h_j(\tilde{x}) - h_j(x^*) \geq \Phi(\tilde{x}, x^*, (\nabla h_j(x^*) \rho_j^*)), j = 1, \dots, p. \quad (14)$$

By (12) – (14), the convexity of $\Phi(x, x^*, \cdot)$ on \mathbb{R}^{n+1} , and the fact that $\tilde{x} \in X$, it

follows that

$$\begin{aligned}
& \Phi\left(\tilde{x}, x^*, \frac{1}{\gamma}\left(\sum_{i=1}^s t_i^* \{\nabla f(x^*, \bar{y}_i) + Au - k_0(\nabla g(x^*, \bar{y}_i) - B\nu)\} + \right.\right. \\
& \quad \left.\left. \sum_{j=1}^p \mu_j^* \nabla h_j(x^*), \sum_{i=1}^s t_i^* (\rho_i + k_0 \bar{\rho}_i) + \sum_{j=1}^p \mu_j^* \rho_j^*\right)\right) \\
& \leq \frac{1}{\gamma} \left[\sum_{i=1}^s t_i^* \Phi(\tilde{x}, x^*, (\nabla f(x^*, \bar{y}_i) + Au, \rho_i)) + k_0 \sum_{i=1}^s t_i^* \Phi(\tilde{x}, x^*, \right. \\
& \quad \left. (-\nabla g(x^*, \bar{y}_i) + B\nu, \bar{\rho}_i)) + \sum_{j=1}^p \mu_j^* \Phi(\tilde{x}, x^*, (\nabla h_j(x^*), \rho_j^*)) \right] \\
& \leq \frac{1}{\gamma} \left[\sum_{i=1}^s t_i^* ((f(\tilde{x}, \bar{y}_i) + \langle \tilde{x}, Au \rangle) - (f(x^*, \bar{y}_i) + \langle x^*, Au \rangle)) + \sum_{i=1}^s t_i^* k_0 \right. \\
& \quad \left. \left(- (g(\tilde{x}, \bar{y}_i) - \langle \tilde{x}, B\nu \rangle) + (g(x^*, \bar{y}_i) - \langle x^*, B\nu \rangle) \right) + \right. \\
& \quad \left. \sum_{j=1}^p \mu_j^* (h_j(\tilde{x}) - h_j(x^*)) \right] < 0, \tag{15}
\end{aligned}$$

where

$$\gamma = 1 + k_0 + \sum_{j=1}^p \mu_j^*.$$

By (3), (9) and the definition of Φ , we have

$$\begin{aligned}
& \Phi\left(\tilde{x}, x^*, \frac{1}{\gamma}\left(\sum_{i=1}^s t_i^* \{\nabla f(x^*, \bar{y}_i) + Au - k_0(\nabla g(x^*, \bar{y}_i) - B\nu)\} \right.\right. \\
& \quad \left.\left. + \sum_{j=1}^p \mu_j^* \nabla h_j(x^*), \sum_{i=1}^s t_i^* (\rho_i + k_0 \bar{\rho}_i) + \sum_{j=1}^p \mu_j^* \rho_j^*\right)\right) \geq 0,
\end{aligned}$$

which is a contradiction to (15).

This completes the proof. \square

4. FIRST DUAL MODEL

In this section, for the considered nondifferentiable generalized minimax fractional programming problem (P), we formulate the first dual model (DI). Further, under (Φ, ρ) -invexity hypotheses, we establish various duality results between (P) and (DI).

Related to (P), we formulate the first dual model (DI) as follows:

$$(DI) \quad \max_{(s,t,\bar{y}) \in K(z)} \sup_{(z,\mu,k,u,\nu) \in \Omega_1(s,t,\bar{y})} k,$$

where $\Omega_1(s,t,\bar{y})$ denotes the set of all $(z,\mu,k,u,\nu) \in \mathbb{R}^n \times \mathbb{R}_+^p \times \mathbb{R}_+ \times \mathbb{R}^n \times \mathbb{R}^n$ satisfying

$$\sum_{i=1}^s t_i \{ \nabla f(z, \bar{y}_i) + Au - k(\nabla g(z, \bar{y}_i) - B\nu) \} + \nabla \langle \mu, h(z) \rangle = 0, \quad (16)$$

$$\sum_{i=1}^s \nabla f(z, \bar{y}_i) + \langle z, Au \rangle - k(g(z, \bar{y}_i) - \langle z, B\nu \rangle) \geq 0, \quad (17)$$

$$\langle \mu, h(z) \rangle \geq 0, \quad (18)$$

$$(s, t, \bar{y}) \in K(z), \quad (19)$$

$$\langle u, Au \rangle \leq 1, \langle \nu, B\nu \rangle \leq 1. \quad (20)$$

For a triplet $(s, t, \bar{y}) \in K(z)$, if the set $\Omega_1(s, t, \bar{y}) = \emptyset$, then we define the supremum over it to be $-\infty$. Throughout this section, we denote

$$\phi_1(\cdot) = \sum_{i=1}^s t_i \{ f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle - k(g(\cdot, \bar{y}_i) - \langle \cdot, B\nu \rangle) \}.$$

Further, let $\text{pr}_{\mathbb{R}^n} \Omega_1(s, t, \bar{y})$ denote the projection of the set $\Omega_1(s, t, \bar{y})$ on \mathbb{R}^n .

Now, we derive weak, strong and strict converse duality theorems.

Theorem 8. (Weak duality) *Let x and $(z, \mu, k, u, \nu, s, t, \bar{y})$ be any feasible solutions of (P) and (DI), respectively. Further, assume that $f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle$, $i = 1, \dots, s$ are (Φ, ρ_i) -invex at z on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_1(s, t, \bar{y})$, $-g(\cdot, \bar{y}_i) + \langle \cdot, B\nu \rangle$, $i = 1, \dots, s$ are $(\Phi, \bar{\rho}_i)$ -invex at z on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_1(s, t, \bar{y})$, $h_j(\cdot)$, $j = 1, \dots, p$ are (Φ, ρ_j^*) -invex at z on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_1(s, t, \bar{y})$ and the inequality*

$$\sum_{i=1}^s t_i (\rho_i + k\bar{\rho}_i) + \sum_{j=1}^p \mu_j \rho_j^* \geq 0, \quad (21)$$

holds. Then,

$$\sup_{y \in Y} \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} \geq k.$$

Proof. We proceed by contradiction. Suppose that

$$\sup_{y \in Y} \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} < k.$$

Therefore, we get the following inequality

$$f(x, \bar{y}_i) + \langle x, Ax \rangle^{1/2} - k \left(g(x, \bar{y}_i) - \langle x, Bx \rangle^{1/2} \right) < 0, \text{ for all } \bar{y}_i \in Y.$$

From $t_i \geq 0, i = 1, \dots, s$ with $\sum_{i=1}^s t_i = 1$, it follows that

$$t_i \left\{ f(x, \bar{y}_i) + \langle x, Ax \rangle^{1/2} - k \left(g(x, \bar{y}_i) - \langle x, Bx \rangle^{1/2} \right) \right\} \leq 0, \quad (22)$$

with at least one strict inequality, because $t = (t_1, \dots, t_s) \neq 0$.

From (1), (16), (19) and (22), we have

$$\begin{aligned} \phi_1(x) &= \sum_{i=1}^s t_i \{ f(x, \bar{y}_i) + \langle x, Au \rangle - k (g(x, \bar{y}_i) - \langle x, Bv \rangle) \} \\ &\leq \sum_{i=1}^s t_i \left\{ f(x, \bar{y}_i) + \langle x, Ax \rangle^{1/2} - k \left(g(x, \bar{y}_i) - \langle x, Bx \rangle^{1/2} \right) \right\} \\ &< 0 \leq \sum_{i=1}^s t_i \{ f(z, \bar{y}_i) + \langle z, Au \rangle - k (g(z, \bar{y}_i) - \langle z, Bv \rangle) \} = \phi_1(z). \end{aligned}$$

Hence,

$$\begin{aligned} &\sum_{i=1}^s t_i \{ (f(x, \bar{y}_i) + \langle x, Au \rangle) - (f(z, \bar{y}_i) + \langle z, Au \rangle) \} \\ &\quad + \sum_{i=1}^s t_i k \{ -(g(x, \bar{y}_i) - \langle x, Bv \rangle) + (g(z, \bar{y}_i) - \langle z, Bv \rangle) \} < 0. \end{aligned}$$

By the feasibility of x for (P) and (18), we get

$$\begin{aligned} &\sum_{i=1}^s t_i \{ (f(x, \bar{y}_i) + \langle x, Au \rangle) - (f(z, \bar{y}_i) + \langle z, Au \rangle) \} + \sum_{i=1}^s t_i k \{ -(g(x, \bar{y}_i) - \langle x, Bv \rangle) \\ &\quad + (g(z, \bar{y}_i) - \langle z, Bv \rangle) \} + \sum_{j=1}^p \mu_j (h_j(x) - h_j(z)) < 0. \quad (23) \end{aligned}$$

From (23) and convexity of $\Phi(x, z, \cdot)$ on \mathbb{R}^{n+1} , it follows that

$$\begin{aligned}
& \Phi\left(x, z, \frac{1}{\gamma}\left(\sum_{i=1}^s t_i\{\nabla f(z, \bar{y}_i) + Au - k(\nabla f(z, \bar{y}_i) - B\nu)\} + \sum_{j=1}^p \mu_j \nabla h_j(z), \right.\right. \\
& \qquad \qquad \qquad \left.\left. \sum_{i=1}^s t_i(\rho_i + k\bar{\rho}_i) + \sum_{j=1}^p \mu_j \rho_j^*\right)\right) \\
& \leq \frac{1}{\gamma}\left(\sum_{i=1}^s t_i \Phi(x, z, (\nabla f(z, \bar{y}_i) + Au, \rho_i)) + k \sum_{i=1}^s t_i \Phi(x, z, (-\nabla g(z, \bar{y}_i) + B\nu, \bar{\rho}_i))\right. \\
& \qquad \qquad \qquad \left. + \sum_{j=1}^p \mu_j \Phi(x, z, (\nabla h_j(z), \rho_j^*))\right) \\
& \leq \frac{1}{\gamma}\left[\sum_{i=1}^s t_i((f(x, \bar{y}_i) + \langle x, Au \rangle) - (f(z, \bar{y}_i) + \langle z, Au \rangle)) + \sum_{i=1}^s t_i k\left(-\langle g(x, \bar{y}_i) \right.\right. \\
& \qquad \qquad \qquad \left.\left. - \langle x, B\nu \rangle) + (g(z, \bar{y}_i) - \langle z, B\nu \rangle)\right) + \sum_{j=1}^p \mu_j^*(h_j(x) - h_j(z))\right] < 0,
\end{aligned} \tag{24}$$

where

$$\gamma = 1 + k + \sum_{j=1}^p \mu_j.$$

By (3), (9) and the definition of Φ we get

$$\begin{aligned}
& \Phi\left(x, z, \frac{1}{\gamma}\left(\sum_{i=1}^s t_i\{\nabla f(z, \bar{y}_i) + Au - k(\nabla f(z, \bar{y}_i) - B\nu)\} + \sum_{j=1}^p \mu_j \nabla h_j(z), \right.\right. \\
& \qquad \qquad \qquad \left.\left. \sum_{i=1}^s t_i(\rho_i + k\bar{\rho}_i) + \sum_{j=1}^p \mu_j \rho_j^*\right)\right) \geq 0,
\end{aligned}$$

which is a contradiction to (24). This completes the proof. \square

Theorem 9. (Strong duality) Assume that x^* is an optimal solution for (P) and that $\nabla h_j(x^*), j \in J(x^*)$, are linearly independent. Then, there exist $(\bar{s}, \bar{t}, \bar{y}^*) \in K(x^*)$ and $(x^*, \bar{\mu}, \bar{k}, \bar{u}, \bar{\nu}) \in \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$ such that $(x^*, \bar{\mu}, \bar{k}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ is a feasible solution for (DI). Further, if all hypotheses of weak duality (Theorem 8) are fulfilled, then $(x^*, \bar{\mu}, \bar{k}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ is optimal in dual problem (DI) and objective functions in problems (P) and (DI) have the same values.

Proof. Since x^* is an optimal solution in the nondifferentiable minimax fractional programming problem (P), by Theorem 6, there exist $(\bar{s}, \bar{t}, \bar{y}^*) \in K(x^*)$ and $(x^*, \bar{\mu}, \bar{k}, \bar{u}, \bar{\nu}) \in \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$ such that $(x^*, \bar{\mu}, \bar{k}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ is feasible for (DI) and

$$\bar{k} = \frac{f(x^*, y_i^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, y_i^*) - \langle x^*, Bx^* \rangle^{1/2}}.$$

Thus, if all the hypotheses of the weak duality (Theorem 8) are fulfilled, then the optimality of $(x^*, \bar{\mu}, \bar{k}, \bar{u}, \bar{v}, \bar{s}, \bar{t}, \bar{y}^*)$ follows directly from this theorem. \square

Theorem 10. (Strict converse duality) Let x^* and $(\bar{z}, \bar{\mu}, \bar{k}, \bar{u}, \bar{v}, \bar{s}, \bar{t}, \bar{y}^*)$ be the optimal solutions for (P) and (DI), respectively, and $\nabla h_j(x^*), j \in J(x^*)$ be linearly independent. Assume that $f(\cdot, y_i^*) + \langle \cdot, A\bar{u} \rangle$ and $-g(\cdot, y_i^*) + \langle \cdot, B\bar{v} \rangle, i = 1, \dots, s$, are strictly (Φ, ρ_i) -invex and strictly $(\Phi, \bar{\rho}_i)$ -invex functions at \bar{z} on $X \cup_{\text{pr}_{\mathbb{R}^n}} \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$ respectively and $h_j(\cdot), j = 1, \dots, p$, is (Φ, ρ_j^*) -invex at \bar{z} on $X \cup_{\text{pr}_{\mathbb{R}^n}} \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$ for all $(\bar{s}, \bar{t}, \bar{y}^*) \in K(x^*)$ and $(\bar{z}, \bar{\mu}, \bar{k}, \bar{u}, \bar{v}) \in \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$. Further, assume that the inequality

$$\sum_{i=1}^s \bar{t}_i(\rho_i + \bar{k}\bar{\rho}_i) + \sum_{j=1}^p \bar{\mu}_j \rho_j^* \geq 0 \quad (25)$$

holds. Then, $x^* = \bar{z}$, that is, \bar{z} is optimal for (P) and

$$\sup_{y \in Y} \frac{f(\bar{z}, \bar{y}^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}}{g(\bar{z}, \bar{y}^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}} = \bar{k}.$$

Proof. Suppose, contrary to the result, that $x^* \neq \bar{z}$. By the hypotheses of the theorem and following along the lines of the proof of Theorem 8, we have

$$\begin{aligned} 0 &\leq \Phi\left(x^*, \bar{z}, \frac{1}{\gamma} \left(\sum_{i=1}^s \bar{t}_i \{ \nabla f(\bar{z}, \bar{y}_i^*) + A\bar{u} - \bar{k}(\nabla g(\bar{z}, \bar{y}_i^*) - B\bar{v}) \} + \sum_{j=1}^p \bar{\mu}_j \nabla h_j(\bar{z}), \right. \right. \\ &\quad \left. \left. \sum_{i=1}^s \bar{t}_i(\rho_i + \bar{k}\bar{\rho}_i) + \sum_{j=1}^p \bar{\mu}_j \rho_j^* \right) \right) \\ &\leq \frac{1}{\gamma} \left(\sum_{i=1}^s \bar{t}_i \Phi(x^*, \bar{z}, (\nabla f(\bar{z}, \bar{y}_i^*) + A\bar{u}, \rho_i)) + \bar{k} \sum_{i=1}^s \bar{t}_i \Phi(x^*, \bar{z}, (-\nabla g(\bar{z}, \bar{y}_i^*) \right. \\ &\quad \left. + B\bar{v}, \bar{\rho}_i)) + \sum_{j=1}^p \bar{\mu}_j \Phi(x^*, \bar{z}, (\nabla h_j(\bar{z}), \rho_j^*)) \right) \\ &< \frac{1}{\gamma} \left[\sum_{i=1}^s \bar{t}_i \{ (f(x^*, \bar{y}_i^*) + \langle x^*, A\bar{u} \rangle) - (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{u} \rangle) \} + \sum_{i=1}^s \bar{t}_i \bar{k} \right. \\ &\quad \left. \{ -(g(x^*, \bar{y}_i^*) - \langle x^*, B\bar{v} \rangle) + (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{v} \rangle) \} + \sum_{j=1}^p \bar{\mu}_j (h_j(x^*) - h_j(\bar{z})) \right], \end{aligned} \quad (26)$$

where

$$\gamma = 1 + \bar{k} + \sum_{j=1}^p \bar{\mu}_j.$$

By the feasibility of x^* and (18), we get

$$\sum_{j=1}^p \bar{\mu}_j [h_j(x^*) - h_j(\bar{z})] \leq 0. \quad (27)$$

From (26) and (27), we have

$$\begin{aligned} \sum_{i=1}^s \bar{t}_i \{ (f(x^*, \bar{y}_i^*) + \langle x^*, A\bar{u} \rangle) - (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{u} \rangle) \} + \\ \sum_{i=1}^s \bar{t}_i \bar{k} \{ -(g(x^*, \bar{y}_i^*) - \langle x^*, B\bar{v} \rangle) + (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{v} \rangle) \} > 0. \end{aligned}$$

By (16) and (2), the above inequality implies that

$$\begin{aligned} \sum_{i=1}^s \bar{t}_i \left\{ \left(f(x^*, \bar{y}_i^*) + \langle x^*, Ax^* \rangle^{1/2} \right) - \bar{k} \left(g(x^*, \bar{y}_i^*) + \langle x^*, Bx^* \rangle^{1/2} \right) \right\} \\ > \sum_{i=1}^s \bar{t}_i \left\{ \left(f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2} \right) - \bar{k} \left(g(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, B\bar{z} \rangle^{1/2} \right) \right\} \geq 0. \end{aligned}$$

From the above inequality, we conclude that there exists a certain $i_0 \in \{1, \dots, s\}$, such that

$$\left\{ \left(f(x^*, \bar{y}_{i_0}^*) + \langle x^*, Ax^* \rangle^{1/2} \right) - \bar{k} \left(g(x^*, \bar{y}_{i_0}^*) + \langle x^*, Bx^* \rangle^{1/2} \right) \right\} > 0.$$

It follows that

$$\sup_{y \in Y} \frac{f(x^*, \bar{y}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x, \bar{y}^*) - \langle x^*, Bx^* \rangle^{1/2}} \geq \frac{f(x^*, \bar{y}_{i_0}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x, \bar{y}^*) - \langle x^*, Bx^* \rangle^{1/2}} > \bar{k}. \quad (28)$$

By the strong duality theorem (Theorem 9), it follows that

$$\sup_{y \in Y} \frac{f(x^*, \bar{y}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x, \bar{y}^*) - \langle x^*, Bx^* \rangle^{1/2}} = \bar{k},$$

which contradicts (28). Therefore, $x^* = \bar{z}$. Hence, the proof is complete. \square

5. SECOND DUAL MODEL

In this section, for the considered nondifferentiable generalized minimax fractional programming problem (P), we formulate the second dual model (DII). Further, under (Φ, ρ) -invexity hypotheses, we establish various duality results between (P) and (DII).

We state the modified version of Theorem 6, by replacing the parameter k_0 with $\frac{f(x^*, \bar{y}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x, \bar{y}^*) - \langle x^*, Bx^* \rangle^{1/2}}$ and by rewriting the multiplier functions associated with the inequality constraints.

Theorem 11. *Let x^* be an optimal solution for (P) and let $\nabla h_j(x^*), j \in J(x^*)$ be linearly independent. Then, there exist $(\bar{s}, \bar{t}, \bar{y}) \in K(x^*)$ and $\bar{\mu} \in \mathbb{R}_+^p$, such that*

$$\begin{aligned} & \sum_{i=1}^{\bar{s}} \bar{t}_i (g(x^*, \bar{y}_i) - \langle x^*, Bx^* \rangle^{1/2}) (\nabla f(x^*, \bar{y}_i) + Au) - (f(x^*, \bar{y}_i) + \\ & \quad \langle x^*, Ax^* \rangle^{1/2}) (\nabla g(x^*, \bar{y}_i) - B\nu) + \sum_{j=1}^p \bar{\mu}_j \nabla h_j(x^*) = 0, \\ & \quad \sum_{i=1}^p \bar{\mu}_i \nabla h_i(x^*) \geq 0, \\ & \quad \bar{\mu} \in \mathbb{R}_+^p, \bar{t}_i \geq 0, \sum_{i=1}^{\bar{s}} \bar{t}_i = 1, \bar{y}_i \in Y(\bar{x}), i = 1, \dots, \bar{s}. \end{aligned}$$

We formulate the second dual model as follows:

$$(DII) \quad \max_{(s,t,\bar{y}) \in K(z)} \sup_{(z,\mu,u,\nu) \in \Omega_2(s,t,\bar{y})} F(z),$$

where $F(z) = \sup_{y \in Y} \frac{f(z,y) + \langle z, Az \rangle^{1/2}}{g(z,y) - \langle z, Bz \rangle^{1/2}}$ and $\Omega_2(s, t, \bar{y})$ denote the set of all $(z, \mu, u, \nu) \in \mathbb{R}^n \times \mathbb{R}_+^m \times \mathbb{R}^n \times \mathbb{R}^n$ satisfying

$$\begin{aligned} & \sum_{i=1}^s t_i \left\{ (g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2}) (\nabla f(z, \bar{y}_i) + Au) - \right. \\ & \quad \left. (f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2}) (\nabla g(z, \bar{y}_i) - B\nu) \right\} + \sum_{j=1}^p \bar{\mu}_j \nabla h_j(z) = 0, \end{aligned} \quad (29)$$

$$\sum_{j=1}^p \mu_j h_j(z) \geq 0, \quad (30)$$

$$(s, t, \bar{y}) \in K(z), \quad (31)$$

$$\langle z, Az \rangle^{1/2} = \langle z, Au \rangle, \langle z, Bz \rangle^{1/2} = \langle z, B\nu \rangle, \langle u, Au \rangle \leq 1, \langle \nu, B\nu \rangle \leq 1 \quad (32)$$

For a triplet $(s, t, \bar{y}) \in K(z)$, if the set $\Omega_2(s, t, \bar{y})$ is empty, then we define the supremum over it to be $-\infty$. Throughout this section, we denote

$$\begin{aligned} \phi_2(\cdot) = & \sum_{i=1}^s t_i \left\{ (g(z, \bar{y}_i) - \langle z, B\nu \rangle) (f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle) - \right. \\ & \left. (f(z, \bar{y}_i) + \langle z, Au \rangle) (\nabla g(\cdot, \bar{y}_i) - \langle \cdot, B\nu \rangle) \right\}. \end{aligned}$$

Further, let $\text{pr}_{\mathbb{R}^n} \Omega_2(s, t, \bar{y})$ denote the projection of the set on \mathbb{R}^n .

Now, we establish weak, strong and strict converse duality theorems.

Theorem 12. (Weak duality) Let x and $(z, \mu, u, \nu, s, t, \bar{y})$ be the feasible solutions for (P) and (DII), respectively. Further, assume that $f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle$, $i = 1, \dots, s$ are (Φ, ρ_i) -invex at z on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_2(s, t, \bar{y})$, $-g(\cdot, \bar{y}_i) + \langle \cdot, B\nu \rangle$, $i = 1, \dots, s$, are $(\Phi, \bar{\rho}_i)$ -invex at z on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_2(s, t, \bar{y})$, $h_j(\cdot)$, $j = 1, \dots, p$ are (Φ, ρ_j^*) -invex at z on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_2(s, t, \bar{y})$ and the inequality

$$\sum_{i=1}^s t_i \left\{ \left(g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) \rho_i + \left(\left(f(z, \bar{y}_i) + \langle z, Bz \rangle^{1/2} \right) \bar{\rho}_i \right) \right\} + \sum_{j=1}^p \mu_j \rho_j^* \geq 0, \quad (33)$$

holds. Then,

$$\sup_{y \in Y} \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} \geq F(z).$$

Proof. Suppose, contrary to the result, that

$$\sup_{y \in Y} \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} < F(z). \quad (34)$$

Since, $\bar{y}_i \in Y(z)$, $i = 1, \dots, s$ we get

$$F(z) = \sup_{y \in Y} \frac{f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2}}{g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2}}. \quad (35)$$

By (34) and (35), we get

$$\begin{aligned} & \left(g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) \left(f(x, \bar{y}_i) + \langle x, Ax \rangle^{1/2} \right) - \\ & \left(f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2} \right) \left(g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) < 0, \end{aligned}$$

for all $i = 1, \dots, s$ and $\bar{y}_i \in Y$.

From $\bar{y}_i \in Y(z) \subset Y$ and $t \in \mathbb{R}_+^s$ with $\sum_{i=1}^s t_i = 1$, it follows that

$$\begin{aligned} & \sum_{i=1}^s t_i \left\{ \left(g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) \left(f(x, \bar{y}_i) + \langle x, Ax \rangle^{1/2} \right) - \right. \\ & \left. \left(f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2} \right) \left(g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) \right\} < 0. \end{aligned} \quad (36)$$

From (1), (32) and (36), we have

$$\begin{aligned} \phi_2(x) &= \sum_{i=1}^s t_i \left\{ (g(z, \bar{y}_i) - \langle z, B\nu \rangle)(f(x, \bar{y}_i) + \langle x, Au \rangle) - \right. \\ &\quad \left. (f(z, \bar{y}_i) + \langle z, Au \rangle)(g(z, \bar{y}_i) - \langle x, Bx \rangle) \right\} \\ &\leq \sum_{i=1}^s t_i \left\{ (g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2})(f(x, \bar{y}_i) + \langle x, Ax \rangle^{1/2}) \right. \\ &\quad \left. - (f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2})(g(x, \bar{y}_i) - \langle x, Bx \rangle^{1/2}) \right\} < 0 = \phi_2(z). \end{aligned}$$

Hence,

$$\phi_2(x) < \phi_2(z).$$

By the feasibility of x for (P) and (30), we get

$$(\phi_2(x) - \phi_2(z)) + \sum_{j=1}^p \mu_j (h_j(x) - h_j(z)) < 0. \quad (37)$$

From the (Φ, ρ_i) -invexity of $(f(\cdot, \bar{y}_i) + \langle \cdot, Au \rangle)$ and $(\Phi, \bar{\rho}_i)$ -invexity of $(-g(\cdot, \bar{y}_i) + \langle \cdot, B\nu \rangle)$ at z , for $i = 1, \dots, s$, it follows that

$$\begin{aligned} f(x, \bar{y}_i) + \langle x, Au \rangle - f(z, \bar{y}_i) - \langle z, Au \rangle &\geq \\ \Phi(x, z, (\nabla f(z, \bar{y}_i) + Au, \rho_i)), &i = 1, \dots, s, \quad (38) \end{aligned}$$

and

$$\begin{aligned} -g(x, \bar{y}_i) + \langle x, B\nu \rangle + g(z, \bar{y}_i) - \langle z, B\nu \rangle &\geq \\ \Phi(x, z, (-\nabla g(z, \bar{y}_i) + B\nu, \bar{\rho}_i)), &i = 1, \dots, s. \quad (39) \end{aligned}$$

The (Φ, ρ_j^*) -invexity of $h_j(\cdot)$ at z implies that

$$h_j(x) - h_j(z) \geq \Phi(x, z, (\nabla h_j(z), \rho_j^*)), j = 1, \dots, p. \quad (40)$$

Multiplying (38) by $t_i (g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2})$, (39) by $t_i (f(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2})$, and then summing up these inequalities to (40) along with the convexity of $\Phi(x, z, (\cdot))$

on \mathbb{R}^{n+1} , we get

$$\begin{aligned}
& \frac{1}{\gamma}(\phi_2(x) - \phi_2(z) + \sum_{j=1}^p \mu_j h_j(x) - \sum_{j=1}^p \mu_j h_j(z)) \\
& \geq \frac{1}{\gamma} \sum_{i=1}^s t_i \left\{ \left(\Phi(x, z, (\nabla f(z, \bar{y}_i) + Au, \rho_i)) \right) (g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2}) \right. \\
& \quad \left. + \left(\Phi(x, z, (\nabla g(z) - B\nu, \bar{\rho}_i)) \right) \left(f(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) \right\} \\
& \quad \quad \quad + \frac{1}{\gamma} \sum_{j=1}^p \mu_j \Phi(x, z, (\nabla h_j(z), \bar{\rho}_j^*)) \\
& \geq \Phi \left(x, z, \frac{1}{\gamma} \left(\sum_{i=1}^s t_i \left(\left(\nabla f(z, \bar{y}_i) + Au \right) (g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2}) \right. \right. \right. \\
& \quad \left. \left. \left. - (f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2}) (\nabla g(z, \bar{y}_i) - B\nu) \right) + \sum_{j=1}^p \mu_j \nabla h_j(z) \right), \frac{1}{\gamma} \left(\rho_i (f(z, \bar{y}_i) \right. \right. \right. \\
& \quad \left. \left. \left. - \langle z, Bz \rangle^{1/2}) + \bar{\rho}_i (f(z, \bar{y}_i) + \langle z, Az \rangle^{1/2}) \right) + \sum_{j=1}^p \bar{\mu}_j \rho_j^* \right) \geq 0,
\end{aligned} \tag{41}$$

where

$$\gamma = \sum_{i=1}^s \bar{t}_i \left\{ \left(g(z, \bar{y}_i) - \langle z, Bz \rangle^{1/2} \right) + \left(f(z, \bar{y}_i) + \langle z, Bz \rangle^{1/2} \right) \right\} + \sum_{j=1}^p \bar{\mu}_j.$$

which is a contradiction to (37). This completes the proof. \square

Theorem 13. (Strong duality) Let x^* be an optimal solution for (P) and $\nabla h_j(x^*)$, $j \in J(x^*)$ be linearly independent. Then, there exist $(\bar{s}, \bar{t}, \bar{y}^*) \in K(x^*)$ and $(x^*, \bar{\mu}, \bar{u}, \bar{\nu}) \in \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$ such that $(x^*, \bar{\mu}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ is a feasible solution for (DII). Further, if all the hypotheses of the weak duality (Theorem 12) are fulfilled, then, $(x^*, \bar{\mu}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ is optimal in dual problem (DII) and objective functions in problems (P) and (DII) have the same values.

Proof. Since x^* is an optimal solution in the considered nondifferentiable minimax fractional programming problem (P), then by Theorem 6 there exist $\bar{u}, \bar{\nu} \in \mathbb{R}^n$ and $\bar{\mu} \in \mathbb{R}_+^p$ to satisfy the expression (29) obtained by substituting

$$k_0 = \frac{f(x^*, \bar{y}_i^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x, \bar{y}_i^*) - \langle x^*, Bx^* \rangle^{1/2}},$$

in (3). It follows that there exist $(\bar{s}, \bar{t}, \bar{y}^*) \in K(x^*)$ and $(x^*, \bar{\mu}, \bar{u}, \bar{\nu}) \in \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$ such that $(x^*, \bar{\mu}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ is feasible for (DII). Thus, if all the hypotheses of the weak duality (Theorem 12) are fulfilled, then, the optimality of $(x^*, \bar{\mu}, \bar{u}, \bar{\nu}, \bar{s}, \bar{t}, \bar{y}^*)$ follows directly from this theorem. \square

Theorem 14. (Strict converse duality) Let x^* and $(\bar{z}, \bar{\mu}, \bar{k}\bar{u}, \bar{v}, \bar{s}, \bar{t}, \bar{y}^*)$ be the optimal solutions for (P) and (DII), respectively and let $\nabla h_j(x^*), j \in J(x^*)$ be linearly independent. Assume that $f(\cdot, \bar{y}_i^*) + \langle \cdot, A\bar{u} \rangle$ and $-g(\cdot, \bar{y}_i^*) + \langle \cdot, B\bar{v} \rangle$, for $i = 1, \dots, s$ are strictly (Φ, ρ_i) -invex and strictly $(\Phi, \bar{\rho}_i)$ -invex at \bar{z} on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$, respectively and let $h_j(\cdot)$ for $j = 1, \dots, p$ be (Φ, ρ_j^*) -invex at \bar{z} on $X \cup \text{pr}_{\mathbb{R}^n} \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$, for all $(\bar{s}, \bar{t}, \bar{y}^*) \in K(x^*)$ and $(\bar{z}, \bar{\mu}, \bar{k}\bar{u}, \bar{v}) \in \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$. Further, assume that the inequality

$$\sum_{i=1}^s \bar{t}_i \left((g(\bar{z}, \bar{y}_i) - \langle \bar{z}, B\bar{z} \rangle^{1/2}) \rho_i + (f(\bar{z}, \bar{y}_i) + \langle \bar{z}, B\bar{z} \rangle^{1/2}) \bar{\rho}_i \right) + \sum_{j=1}^p \bar{\mu}_j \rho_j^* \geq 0, \quad (42)$$

holds. Then, $x^* = \bar{z}$, that is, \bar{z} is optimal for (P) and

$$\sup_{y \in Y} \frac{f(\bar{z}, \bar{y}^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}}{g(\bar{z}, \bar{y}^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}} = F(\bar{z}).$$

Proof. Suppose, contrary to the result, that $x^* \neq \bar{z}$. Using the hypotheses of the theorem and proceeding as in Theorem 12, we get

$$\begin{aligned} & \frac{1}{\gamma} \left(\phi_2(x^*) - \phi_2(\bar{z}) + \sum_{j=1}^p \bar{\mu}_j h_j(x^*) - \sum_{j=1}^p \bar{\mu}_j h_j(\bar{z}) \right) \\ & > \frac{1}{\gamma} \sum_{i=1}^s \bar{t}_i \left\{ (\Phi(x^*, \bar{z}, (\nabla f(\bar{z}, \bar{y}_i^*) + A\bar{u}, \rho_i))) (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}) \right. \\ & \quad \left. + (\Phi(x^*, \bar{z}, (\nabla g(\bar{z}) + B\bar{v}, \bar{\rho}_i))) (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}) \right\} + \frac{1}{\gamma} \sum_{j=1}^p \bar{\mu}_j \\ & \Phi(x^*, \bar{z}, (\nabla h_j(\bar{z}), \rho_j^*)) \\ & \geq \Phi \left(x^*, \bar{z}, \frac{1}{\gamma} \left(\sum_{i=1}^s \bar{t}_i \left((\nabla f(\bar{z}, \bar{y}_i^*) + A\bar{u}) (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}) \right. \right. \right. \\ & \quad \left. \left. - (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}) (\nabla g(\bar{z}, \bar{y}_i^*) - B\bar{v}) \right) + \sum_{j=1}^p \bar{\mu}_j \nabla h_j(\bar{z}) \right), \\ & \quad \left. \rho_i (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}) + \bar{\rho}_i (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}) + \sum_{j=1}^p \bar{\mu}_j \rho_j^* \right) \geq 0, \end{aligned}$$

where

$$\gamma = \sum_{i=1}^s \bar{t}_i \left\{ (g(\bar{z}, \bar{y}_i) - \langle \bar{z}, B\bar{z} \rangle^{1/2}) + (f(\bar{z}, \bar{y}_i) + \langle \bar{z}, B\bar{z} \rangle^{1/2}) \right\} + \sum_{j=1}^p \bar{\mu}_j.$$

That is,

$$\frac{1}{\gamma} \left(\phi_2(x^*) - \phi_2(\bar{z}) + \sum_{j=1}^p \bar{\mu}_j h_j(x^*) - \sum_{j=1}^p \bar{\mu}_j h_j(\bar{z}) \right) > 0. \quad (43)$$

By the feasibility of x^* and (30), we get

$$\sum_{j=1}^p \bar{\mu}_j [h_j(x^*) - h_j(\bar{z})] \leq 0. \quad (44)$$

From (43) and (44), we have

$$\phi_2(x^*) - \phi_2(\bar{z}) > 0.$$

That is,

$$\begin{aligned} & \sum_{i=1}^{\bar{s}} \bar{t}_i \left\{ (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{v} \rangle)(f(x^*, \bar{y}_i^*) + \langle x^*, A\bar{u} \rangle) - (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{u} \rangle)(g(x^*, \bar{y}_i^*) - \langle x^*, B\bar{v} \rangle) \right\} \\ & > \sum_{i=1}^{\bar{s}} \bar{t}_i \left\{ (g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{v} \rangle)(f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{u} \rangle) - (f(\bar{z}, \bar{y}_i^*) + \langle \bar{z}, A\bar{u} \rangle)(g(\bar{z}, \bar{y}_i^*) - \langle \bar{z}, B\bar{v} \rangle) \right\} \geq 0. \end{aligned}$$

Therefore, there exists a certain i_0 , such that

$$\begin{aligned} & \sum_{i=1}^{\bar{s}} \bar{t}_i \left\{ (g(\bar{z}, \bar{y}_{i_0}^*) - \langle \bar{z}, B\bar{v} \rangle)(f(x^*, \bar{y}_{i_0}^*) + \langle x^*, A\bar{u} \rangle) - \right. \\ & \left. (f(\bar{z}, \bar{y}_{i_0}^*) + \langle \bar{z}, A\bar{u} \rangle)(g(x^*, \bar{y}_{i_0}^*) - \langle x^*, B\bar{v} \rangle) \right\} > 0. \end{aligned}$$

From the above inequality and (32), it follows that

$$\sup_{y \in Y} \frac{f(x^*, \bar{y}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, \bar{y}^*) - \langle x^*, Bx^* \rangle^{1/2}} \geq \frac{f(x^*, \bar{y}_{i_0}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, \bar{y}_{i_0}^*) - \langle x^*, Bx^* \rangle^{1/2}} > F(\bar{z}). \quad (45)$$

By the strong duality theorem (Theorem 13), it follows that

$$\sup_{y \in Y} \frac{f(x^*, \bar{y}^*) + \langle x^*, Ax^* \rangle^{1/2}}{g(x^*, \bar{y}^*) - \langle x^*, Bx^* \rangle^{1/2}} = F(\bar{z}). \quad (46)$$

Thus, inequality (46), contradicts (45). Therefore, $x^* = z$. Hence, the proof is complete. \square

The following example illustrates the significance of Theorems 6-14.

Example 15. Consider the following nondifferentiable generalized minimax fractional programming problem:

$$(P1) \quad \inf_{x \in \mathbb{R}} \sup_{y \in Y} \left\{ \phi(x, y) := \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} \right\}$$

subject to $h(x) \leq 0$,

where $Y = [0, 1]$, $A = B = 1$. The functions $f, g : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and $h : \mathbb{R} \rightarrow \mathbb{R}$ are defined as

$$f(x, y) = x^2 + x + y^2, \quad g(x, y) = 2x + 1,$$

and

$$h(x) = 1 - x.$$

Note that the set of all feasible solutions is $X = \{x \in \mathbb{R} : 1 - x \leq 0\} = [1, \infty)$. The sets $Y(x)$ and $K(x)$ are given as

$$Y(x) = \{1\} \quad \text{and} \quad K(x) = \{(1, 1, 1)\}.$$

Now, for all $(x, y) \in X \times Y = \{(x, y) : x \geq 1, 0 \leq y \leq 1\}$, we have

$$f(x, y) + \langle x, Ax \rangle^{1/2} = x^2 + x + y^2 + |x| > 0$$

and

$$g(x, y) - \langle x, Bx \rangle = 2x + 1 - |x| > 0.$$

Now, we will find a minimax solution of (P) for $x^* \in [1, \infty)$, we have considered the following cases.

Case 1 Let $x^* = 1$. From (3)-(8), we get $x^* = 1, u = 1, v = 1, k_0 = 2$ and $\mu^* = 2$. Hence, necessary optimality conditions (Theorem 6) are satisfied at x^* .

Case 2 Let $x^* > 1$. From (4)-(8), we get $u = 1, v = 1, k_0 = x^* + 1$ and $\mu^* = 0$. Putting these values in (3), we get $x^* + 1 = 0$, which is not possible for any $x^* > 1$. Hence, necessary optimality conditions (Theorem 6) are not satisfied for $x^* > 1$.

Define

$$\Phi(x, y, (z, \rho)) = \langle z, x - y \rangle + \rho \|x - y\|^2.$$

We can check that the functions $f(x, \bar{y}) + \langle x, Au \rangle = x^2 + x + \bar{y}^2 + xu$ and $-g(x, \bar{y}) + \langle x, Bv \rangle = -2x - 1 + xv$ are (Φ, ρ) -invex for $\rho = 1/2$ and $(\Phi, \bar{\rho})$ -invex at x^* for $\bar{\rho} = -1/8$ on X , respectively, and $h(x)$ is (Φ, ρ^*) -invex at x^* for $\rho^* = -1/8$ on X . Furthermore,

$$t^*(\rho + k_0 \bar{\rho}) + \mu^* \rho^* = 0.$$

By setting $s = 1, t^* = 1, \bar{y} = 1, u = 1, v = 1, k_0 = 2$ and $\mu^* = 2$, the sufficient optimality conditions (Theorem 7) are satisfied. Hence, $x^* = 1$ is the optimal solution for (P1).

Now, we will formulate the first dual model for (P1).

$$(DI') \quad \max_{(s, t, \bar{y}) \in K(z)} \sup_{(z, \mu, k, u, v) \in \Omega_1(s, t, \bar{y})} k,$$

where Ω_1 denotes the set of all $(z, \mu, k, u, v) \in \mathbb{R} \times \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R} \times \mathbb{R}$ satisfying

$$\begin{aligned} 2z + 1 + u - k(2 - v) - \mu &= 0, \\ 2z + 1 + zu - k(2z + 1 - zv) &\geq 0 \\ \mu(1 - z) &\geq 0, \\ u^2 \leq 1, v^2 &\leq 1. \end{aligned}$$

We can verify that, $\Omega_1 = \{(z, \mu, k, u, v) : 0 \leq z \leq 1, 0 \leq \mu \leq 4, 0 \leq k \leq 2, -1 \leq u, v \leq 1\}$.

We can check that the functions $f(x, \bar{y}) + \langle x, Au \rangle = x^2 + x + \bar{y}^2 + xu$ and $-g(x, \bar{y}) + \langle x, Bv \rangle = -2x - 1 + xv$ are strictly (Φ, ρ) -invex for $\rho = 1/2$ and strictly $(\Phi, \bar{\rho})$ -invex at x^* for $\bar{\rho} = -1/8$ on $X \cup_{pr\mathbb{R}} \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$, respectively, and $h(x)$ is strictly (Φ, ρ^*) -invex at x^* for $\rho^* = -1/8$ on $X \cup_{pr\mathbb{R}} \Omega_1(\bar{s}, \bar{t}, \bar{y}^*)$. Furthermore,

$$t^*(\rho + k_0\bar{\rho}) + \mu^* \rho^* = 0,$$

and

$$\sup_{y \in Y} \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} \geq k, \quad \forall x \in X \text{ and } k \in \Omega_1(s, t, \bar{y}).$$

Hence, the weak duality conditions (Theorem 8) are satisfied.

By setting $\bar{s} = 1, \bar{t} = 1, \bar{y}^* = 1, \bar{u} = 1, \bar{v} = 1, \bar{k}_0 = 2$ and $\bar{\mu} = 2$, the strong converse duality conditions (Theorem 9) and strict converse duality conditions (Theorem 10) are satisfied. Hence, $\bar{z} = x^* = 1$ is the optimal solution for (P1) and

$$\sup_{y \in Y} \frac{f(\bar{z}, \bar{y}^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}}{g(\bar{z}, \bar{y}^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}} = \bar{k}.$$

Now, we will formulate the second dual model for (P1).

$$(DII') \quad \max_{(s, t, \bar{y}) \in K(z)} \sup_{(z, \mu, u, v) \in \Omega_2(s, t, \bar{y})} F(z),$$

where $F(z) = \sup_{y \in Y} \frac{f(z, y) + \langle z, Az \rangle^{1/2}}{g(z, y) - \langle z, Bz \rangle^{1/2}}$ and Ω_2 denotes the set of all $(z, \mu, u, v) \in \mathbb{R} \times \mathbb{R}_+ \times \mathbb{R} \times \mathbb{R}$ satisfying

$$\begin{aligned} (2z + 1 - |z|)(2z + 1 + u) - (z^2 + z + 1 + |z|)(2 - v) - \mu &= 0, \\ \mu(1 - z) &\geq 0, \\ u^2 \leq 1, v^2 \leq 1, |z| = zu, |z| &= zv. \end{aligned}$$

We can verify that, $\Omega_2 = \{(z, \mu, u, v) : z \in (-\infty, \frac{-1-\sqrt{10}}{3}] \cup [0, 1], \mu \in [1, 4], u = v \in \{-1, 0, 1\}\}$.

We can check that the functions $f(x, \bar{y}) + \langle x, Au \rangle = x^2 + x + \bar{y}^2 + xu$ and $-g(x, \bar{y}) + \langle x, Bv \rangle = -2x - 1 + xv$ are strictly (Φ, ρ) -invex for $\rho = 1/2$ and strictly

$(\Phi, \bar{\rho})$ -invex at x^* for $\bar{\rho} = -1/8$ on $X \cup_{pr_{\mathbb{R}}} \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$, respectively, and $h(x)$ is strictly (Φ, ρ^*) -invex at x^* for $\rho^* = -1/8$ on $X \cup_{pr_{\mathbb{R}}} \Omega_2(\bar{s}, \bar{t}, \bar{y}^*)$. Furthermore,

$$t^* \{ (g(z, \bar{y}) - \langle z, Bz \rangle^{1/2})\rho + (f(z, \bar{y} + \langle z, Bz \rangle^{1/2})\bar{\rho}) \} + \mu\rho^* \geq 0, \forall z \in [0, 1].$$

and

$$\sup_{y \in Y} \frac{f(x, y) + \langle x, Ax \rangle^{1/2}}{g(x, y) - \langle x, Bx \rangle^{1/2}} \geq F(z), \forall x \in X \text{ and } z \in \Omega_2(s, t, \bar{y}).$$

Hence, weak duality conditions (Theorem 12) are satisfied.

By setting $\bar{z} = 1, \bar{s} = 1, \bar{t} = 1, \bar{y}^* = 1, \bar{u} = 1, \bar{v} = 1$ and $\bar{\mu} = 2$, the strong converse duality conditions (Theorem 13) and strict converse duality conditions (Theorem 14) are satisfied. Hence, $\bar{z} = x^* = 1$ is the optimal solution for (P1) and

$$\sup_{y \in Y} \frac{f(\bar{z}, \bar{y}^*) + \langle \bar{z}, A\bar{z} \rangle^{1/2}}{g(\bar{z}, \bar{y}^*) - \langle \bar{z}, B\bar{z} \rangle^{1/2}} = F(\bar{z}).$$

6. CONCLUSIONS and FUTURE DIRECTIONS

In the paper, a new class of nonconvex nondifferentiable generalized minmax programming problems (P) has been considered. The sufficient optimality results have been established for such nonsmooth optimization problems under hypotheses that the functions involved are (Φ, ρ) -invex. Further, two dual models (DI) and (DII) have been formulated for the considered nondifferentiable generalized minmax programming problem (P) and several duality results have been established between the primal optimization problem and its duals also under (Φ, ρ) -invexity. Note that, in the light of Remark 2.5, the results of the paper extend and generalize several results of Lai and Lee [29], Lai et al. [30], Liang and Shi [32], Liu and Wu [33], Liu et al. [35] and Mishra et al. [37, 40]. Results of the paper may be extended for real valued nondifferentiable locally Lipschitz functions on real Banach spaces using the tools of Michel Penot subdifferentials or convexificators, which will orient the future research of the authors.

REFERENCES

- [1] Antczak, T., “Generalized fractional minimax programming with B - (p, r) -invexity”, *Computers & Mathematics with Applications*, 56 (6) (2008) 1505–1525.
- [2] Antczak, T., “Nonsmooth minimax programming under locally Lipschitz (Φ, ρ) -invexity”, *Applied Mathematics and Computation*, 217 (1) (2011) 9606–9624.
- [3] Antczak, T., Mishra, S. K., and Upadhyay, B. B., “First order duality results for a new class of nonconvex semi-infinite minimax fractional programming problems”, *Journal of Advanced Mathematical Studies*, 9 (1) (2016) 132–162.
- [4] Antczak, T., Mishra, S. K., and Upadhyay, B. B., “Optimality conditions and duality for generalized fractional minimax program involving locally Lipschitz (b, ψ, Φ, ρ) -invexity”, *Control and Cybernetics*, 47 (1) (2018) 5–32.

- [5] Antczak, T., and Singh, V., “Optimality and duality for minimax fractional programming with support functions under $B-(p,r)$ -Type I assumptions”, *Mathematical and Computer Modelling*, 57 (5-6) (2013) 1083–1100.
- [6] Antczak, T., and Stasiak, A., “ (Φ, ρ) -invexity in nonsmooth optimization”, *Numerical Functional Analysis and Optimization*, 32 (1) (2011) 1–25.
- [7] Bajona-Xandri, C., *On Fractional Programming*, Masters Thesis, Universitat Autònoma de Barcelona, 1993.
- [8] Barrodale, I., “Best rational approximation and strict-quasiconvexity”, *SIAM Journal on Numerical Analysis*, 10 (1) (1973) 8–12.
- [9] Bector, C. R., and Bhatia, B. L., “Sufficient optimality and duality for minimax problem”, *Utilitas Mathematica*, 27 (1985) 229–247.
- [10] Bector, C. R., Chandra, S., and Kumar, V., “Duality for minimax programming involving V -invex functions”, *Optimization*, 30 (2) (1994) 93–103.
- [11] Bram, J., “The Lagrange multiplier theorem for max-min with several constraints”, *SIAM Journal on Applied Mathematics*, 14 (4) (1966) 665–667.
- [12] Caristi, G., Ferrara, M., and Stefanescu, A., “*Mathematical programming with (Φ, ρ) -invexity*”, , in: (I.V. Konov, D.T. Luc, A.M. Rubinov), (ed.) *Generalized Convexity and Related Topics: Lecture Notes in Economics and Mathematical Systems* Springer, Berlin, 2006.
- [13] Chandra, S., and Kumar, V., “Duality in fractional minimax programming”, *Journal of the Australian Mathematical Society (series A)*, 58 (3) (1995) 376–386.
- [14] Craven, B. D., “Invex functions and constrained local minima”, *Bulletin of the Australian Mathematical Society*, 24 (3) (1981) 357–366.
- [15] Danskin, J. M., “The theory of max-min with applications”, *SIAM Journal on Applied Mathematics*, 14 (4) (1966) 641–644.
- [16] Dubey, R., Mishra, L. N., and Ali, R., “Special class of second-order non-differentiable symmetric duality problems with (G, α_f) -pseudobonvexity assumptions”, *Mathematics*, 7 (8) (2019) 763.
- [17] Dubey, R., and Mishra, V. N., “Symmetric duality results for a nondifferentiable multi-objective programming problem”, *RAIRO Operations Research*, 53 (2) (2019) 539–558.
- [18] Dubey, R., Kumar, A., Ali, R., and Mishra, L. N., “New class of G-Wolfe type symmetric duality model and duality relations under G_f bonvexity over arbitrary cones”, *Journal of Inequalities and Applications*, 30 (1) (2020) 1-16.
- [19] Dubey, R., Mishra, L. N., Luis Manuel Sánchez Ruiz, “Nondifferentiable G-Mond-Weir type multiobjective symmetric fractional problem and their duality theorems under generalized assumptions”, *Symmetry*, 11 (11) (2019) 1348.
- [20] Dubey, R., Mishra, V. N., and Ali, R., “Duality for Unified Higher-Order Minimax Fractional Programming with Support Function under Type-I Assumptions”, *Mathematics*, 7 (11) (2019) 1034.
- [21] Dubey, R., Deepmala, and Mishra, V. N., “Higher-order symmetric duality in nondifferentiable multiobjective fractional programming problem over cone constraints”, *Statistics, Optimization and Information Computing*, 8 (1) (2020) 187–205.
- [22] Ferrara, M., and Stefanescu, M. V., “Optimality conditions and duality in multiobjective programming with (Φ, ρ) -invexity”, *Yugoslav Journal of Operations Research*, 18 (2) (2008) 153–165.
- [23] Flouda, C. A., and Pardalos, P. M., eds, *Encyclopedia of Optimization*, 2nd ed Springer, New York, 2009.
- [24] Hanson, M. A., “On sufficiency of Kuhn-Tucker conditions”, *Journal of Mathematical Analysis and Applications*, 30 (1981) 545–550.
- [25] Hanson, M. A., and Mond, B., “Further generalizations of convexity in mathematical programming”, *Journal of Information and Optimization Sciences*, 3 (1) (1982) 22–35.
- [26] Husain, I., Hanson, M. A., and Jabeen, Z., “On nondifferentiable minimax fractional programming”, *European Journal of Operational Research*, 160 (2005) 202–217.
- [27] Jayswal, A., “Optimality and duality for nondifferentiable minimax fractional programming with generalized convexity”, *ISRN Applied Mathematics*, 2011, 1-19.
- [28] Jeyakumar, V., “Strong and weak invexity in mathematical programming”, *Methods of*

- Operations Research*, 55 (1995) 109–125.
- [29] Lai, H. C., and Lee, J. C., “On duality theorem for nondifferentiable minimax fractional programming”, *Journal of Computational and Applied Mathematics*, 146 (1) (2002) 115–126.
- [30] Lai, H. C., Liu, J. C., and Tanaka, K., “Necessary and sufficient conditions for minimax fractional programming”, *Journal of Mathematical Analysis and Applications*, 230 (2) (1999) 311–328.
- [31] Liang, Z. A., Huang, H. X., and Pardalos, P. M., “Optimality conditions and duality for nonlinear fractional programming problems”, *Journal of Optimization Theory and Applications*, 110 (2001) 611–619.
- [32] Liang, Z. A., and Shi, J. W., “Optimality conditions and duality for minimax fractional programming with generalized convexity”, *Journal of Mathematical Analysis and Applications*, 277 (2003) 474–488.
- [33] Liu, J. C., and Wu, C. S., “On minimax fractional optimality conditions and (F, ρ) -convexity”, *Journal of Mathematical Analysis and Applications*, 219 (1998) 36–51.
- [34] Liu, J. C., Wu, C. S., and Sheu, R. L., “Duality for fractional minimax programming”, *Optimization*, 41 (2) (2010) 117–133.
- [35] Liu, X. L., Lai, G. M., Xu, C. Q., and Yuan, D. H., “On Nonsmooth semi-infinite minimax programming problem with (Φ, ρ) -invexity”, *Abstract and Applied Analysis*, (3) (2014) 1–7.
- [36] Luo, H. Z., and Wu, H. X., “On necessary conditions for a class of nondifferentiable minimax fractional programming”, *Journal of Computational and Applied Mathematics*, 215 (1) (2008) 103–113.
- [37] Mishra, S. K., Pant, R. P., Rautela, J. S., “Generalized α -invexity and nondifferentiable minimax fractional programming”, *Journal of Computational and Applied Mathematics*, 206 (1) (2007) 122–135.
- [38] Mishra, S. K., and Upadhyay, B. B., “Nonsmooth minimax fractional programming involving η -pseudolinear functions”, *Optimization*, 63 (5) (2014) 775–788.
- [39] Mishra, S. K., and Upadhyay, B. B., *Pseudolinear Functions and Optimization*, Chapman and Hall, CRC Press, UK, 2015.
- [40] Mishra, S. K., Wang, S. Y., Lai, K. K., Shi, J. M., “Nondifferentiable minimax fractional programming under generalized univexity”, *Journal of Computational and Applied Mathematics*, 158 (2) (2003) 379–395.
- [41] Preda, V., “On efficiency and duality for multiobjective programs”, *Journal of Mathematical Analysis and Applications*, 166 (2) (1992) 365–377.
- [42] Schmitendorf, W. E., “Necessary conditions and sufficient conditions for static minimax problems”, *Journal of Mathematical Analysis and Applications*, 57 (1977) 683–693.
- [43] Schroeder, R. G., “Linear programming solutions to ratio games”, *Operational Research*, 18 (1970) 300–305.
- [44] Stefanescu, M. V., Stefanescu, A., “On semi-infinite minimax programming with generalized invexity”, *Optimization*, 61 (11) (2012) 1307–1319.
- [45] Upadhyay, B. B., and Mishra, S. K., “Nonsmooth semi-infinite minimax programming involving generalized (Φ, ρ) -invexity”, *Journal of Systems Science and Complexity*, 28 (4) (2014).
- [46] Vial, J. P., “Strong and weak convexity of sets and functions”, *Mathematics of Operations Research*, 8 (1983) 231–259.
- [47] Weir, T., and Mond, B., “Generalized convexity and duality in multiple objective programming”, *Bulletin of the Australian Mathematical Society*, 39 (2) (1989) 287–299.
- [48] Weir, T., “Pseudoconvex minimax programming”, *Utilitas Mathematica*, 42 (1992) 234–240.
- [49] Yang, X. M., and Hou, S. H., “On minimax fractional optimality and duality with generalized convexity”, *Journal of Global Optimization*, 32 (2005) 235–252.
- [50] Yuan, D. H., Liu, X. L., Chinchuluun, A., and Pardalos, P. M., “Nondifferentiable minimax fractional programming with (C, α, ρ, d) -convexity”, *Journal of Optimization Theory and Applications*, 129 (1) (2006) 185–199.